arXiv:1401.2014v1 [math.AG] 9 Jan 2014

SINGULAR RATIONALLY CONNECTED THREEFOLDS WITH NON-ZERO PLURI-FORMS

WENHAO OU

Abstract. This paper is concerned with singular projective rationally connected threefolds X which carry non-zero pluri-forms, *i.e.* $H^0(X,(\Omega^1_X)^{[\otimes m]})\neq\{0\}$ for some $m>0$, where $(\Omega^1_X)^{[\otimes m]}$ is the reflexive hull of $(\Omega^1_X)^{\otimes m}$. If X has $\mathbb{Q}$-factorial terminal singularities, then we show that there is a fibration p from X to $\mathbb{P}^1$. Moreover, there is a natural isomorphism $H^0(X,(\Omega^1_X)^{[\otimes m]})\cong H^0(\mathbb{P}^1,\mathscr{O}_{\mathbb{P}^1}(-2m+\sum_{z\in\mathbb{P}^1}[\frac{(m(p,z)-1)m}{m(p,z)}]))$ for all $m>0$, where $m(p,z)$ is the smallest positive coefficient in the divisor p^*z.

Contents

1. Introduction 1
2. Preliminaries 2
2.1. Reflexive sheaves on normal varieties 3
2.2. Minimal model program 4
2.3. Examples 5
3. Fibrations over curves 7
3.1. Construction of rationally connected varieties carrying non-zero pluri-forms 7
3.2. The case of threefolds with canonical singularities 8
4. Mori fibrations and non-zero pluri-form 9
4.1. General properties 9
4.2. The case of threefolds with canonical singularities 12
5. Proof of Theorem 1.4 13
References 15

1. Introduction

Recall that a complex projective variety X is said to be rationally connected if for any two general points in X, there exists a rational curve passing through them (see [Kol96, Def. IV.3.2 and Prop. IV.3.6]). It is known that if X is a smooth rationally connected variety, then X does not carry any non-zero pluri-form, *i.e.* $H^0(X,(\Omega^1_X)^{\otimes m})=\{0\}$ for $m>0$ (see [Kol96, Cor. IV.3.8]). It is a conjecture that the converse is also true (see [Kol96, Conj. IV.3.8.1]). For singular varieties, there are also some analogue results. The following theorems can be found in [GKP12] and [GKKP11].

Theorem 1.1 ([GKP12, Thm. 3.3])**.** *If X is a rationally connected variety with factorial canonical singularities, then $H^0(X,(\Omega^1_X)^{[\otimes m]})=\{0\}$ for $m>0$, where $(\Omega^1_X)^{[\otimes m]}$ is the reflexive hull of $(\Omega^1_X)^{\otimes m}$.*

Theorem 1.2 ([GKKP11, Thm. 5.1])**.** *If (X,D) is a klt pair such that X is rationally connected, then $H^0(X,\Omega^{[m]}_X)=\{0\}$ for $m>0$, where $\Omega^{[m]}_X$ is the reflexive hull of Ω^m_X.*

Note that Ω^m_X is a direct summand of $(\Omega^1_X)^{\otimes m}$, hence $H^0(X,(\Omega^1_X)^{[\otimes m]})=\{0\}$ implies $H^0(X,\Omega^{[m]}_X)=\{0\}$. However, there are examples of klt rationally connected varieties whose canonical divisor is effective (see [Tot10, Exmp. 10] or [Kol08, Exmp. 43]). Moreover, Theorem 1.1 is not true if the variety is not factorial. There is a counterexample given in [GKP12, Example 3.7]. In [Ou13], we classify all rationally connected surfaces X with canonical singularities such that $H^0(X,(\Omega^1_X)^{[\otimes m]})\neq\{0\}$ for some $m>0$. We obtain the following theorem.

Theorem 1.3 ([Ou13, Thm. 1.3]). *Let X be a rationally connected surface with canonical singularities. Then $H^0(X,(\Omega^1_X)^{[\otimes m]})\neq\{0\}$ for some $m>0$ if and only if there is a fibration $p:X\to\mathbb{P}^1$ whose general fibers are isomorphic to $\mathbb{P}^1$ such that $\sum_{z\in\mathbb{P}^1}[\frac{m(p,z)-1}{m(p,z)}]\geqslant 2$, where $m(p,z)$ is the smallest positive coefficient in the divisor p^*z.*

In this paper, we will study the case of threefolds. We are interested in the structure of rationally connected threefolds which carry non-zero pluri-forms and we try to find out the source of these forms. If the threefold has terminal singularities, we prove the same result as Theorem 1.3 (see Theorem 1.4 below). In this case, there is a fibration from the threefold to $\mathbb{P}^1$ and the source of non-zero pluri-forms is the ramification of the fibration.

Theorem 1.4. *Let X be a projective rationally connected threefold with $\mathbb{Q}$-factorial terminal singularities. Then $H^0(X,(\Omega^1_X)^{[\otimes m]})\neq\{0\}$ for some $m>0$ if and only if there is a fibration $p:X\to\mathbb{P}^1$ whose general fibers are smooth rationally connected surfaces such that $\sum_{z\in\mathbb{P}^1}\frac{m(p,z)-1}{m(p,z)}\geqslant 2$, where $m(p,z)$ is the smallest positive coefficient in the divisor p^*z. Moreover, if it is the case, then for all $m>0$, we have $H^0(X,(\Omega^1_X)^{[\otimes m]})\cong H^0(\mathbb{P}^1,\mathscr{O}_{\mathbb{P}^1}(-2m+\sum_{z\in\mathbb{P}^1}[\frac{(m(p,z)-1)m}{m(p,z)}]))$, which is of dimension $1-2m+\sum_{z\in\mathbb{P}^1}[\frac{(m(p,z)-1)m}{m(p,z)}]$.*

Construction 1.5. Thanks to Theorem 1.4, every rationally connected threefold X with $\mathbb{Q}$-factorial terminal singularities such that $H^0(X,(\Omega^1_X)^{[\otimes m]})\neq\{0\}$ for some $m>0$ can be constructed as follows. There is a fibration $q:T\to B$ from a normal threefold T to a smooth curve B which has positive genus such that $m(q,b)=1$ for all $b\in B$. There is a finite group G which acts on T and B such that q is G-equivariant. By taking the quotient by G, we have $B/G\cong\mathbb{P}^1$ and $T/G\cong X$. For more details, see §3.

If the threefold X has canonical singularities, we also obtain some necessary conditions for existence of non-zero pluri-forms (see §3.2 and §4). Roughly speaking, if $f_X:X\dashrightarrow X'$ is the result of a MMP for X, then there is a fibration from X' to a variety Z of positive dimension such that the forms come from the base Z. The difficulty in this situation is that the rational map $X\dashrightarrow Z$ is not always regular. And even it is, it may not be equidimensional. If klt singularities are permitted for our threefolds, then there exist other structures. We give an example of rationally connected threefold of general type in Example 2.15.

The main objective of this paper is to proof Theorem 1.4. We will do this in several steps. First, we consider a projective rationally connected threefold X with $\mathbb{Q}$-factorial canonical singularities which carries non-zero pluri-forms. Since it is rationally connected and with canonical singularities, its canonical divisor K_X is not pseudo-effective. Hence if $f_X:X\to X'$ is the result of a MMP for X, then $K_{X'}$ is not pseudo-effective neither. We have a Mori fibration $p:X'\to Z$ where Z is a normal variety of dimension less than 3. However, since X carries non-zero pluri-forms, we can show that $H^0(X',(\Omega^1_{X'})^{[\otimes m]})\neq\{0\}$ for some $m>0$. Then we obtain $\dim Z>0$ by [Ou13, Thm. 2.1]. Hence either $\dim Z=1$ or $\dim Z=2$. We will treat this two cases separately in §3 and §4. If $\dim Z=1$, then $Z\cong\mathbb{P}^1$ and we will show that $K_{\mathbb{P}^1}+\sum_{z\in\mathbb{P}^1}\frac{m(p,z)-1}{m(p,z)}z$ is effective, which is the same condition as in Theorem 1.4. If $\dim Z=2$, then we will define a $\mathbb{Q}$-divisor Δ on Z (see Theorem 4.2 for the definition). We will prove that either $K_Z+\Delta$ is an effective $\mathbb{Q}$-divisor or there is a fibration $X'\to\mathbb{P}^1$ such that we can reduce to the situation of §3. In the last section, we assume that the variety X has terminal singularities. In this case, there is always a fibration from X' to $\mathbb{P}^1$ and we are always in the situation of §3. This fibration induces a dominant rational map from X to $\mathbb{P}^1$. In the end, we will prove that this rational map is regular and complete the proof of Theorem 1.4

2. Preliminaries

Throughout this paper, we will work over $\mathbb{C}$, the field of complex numbers. Unless otherwise specified, every variety is an integral $\mathbb{C}$-scheme of finite type. A curve is a variety of dimension 1, a surface is a variety of dimension 2 and a threefold is a variety of dimension 3. For a normal variety X, let K_X be a canonical divisor of X. We denote the sheaf of Kähler differentials by Ω^1_X. Denote $\bigwedge^p\Omega^1_X$ by Ω^p_X for $p\in\mathbb{N}$. Let $\Omega^{[p]}_X$ (resp. $(\Omega^1_X)^{[\otimes p]}$) be the reflexive hull of Ω^p_X (resp. $(\Omega^1_X)^{\otimes p}$). We say that a normal variety X carries non-zero pluri-forms if $H^0(X,(\Omega^1_X)^{[\otimes m]})\neq\{0\}$ for some $m>0$. Let $\mathrm{Pic}(X)_{\mathbb{Q}}=\mathrm{Pic}(X)\otimes\mathbb{Q}$, where $\mathrm{Pic}(X)$ is the Picard group of X. A $\mathbb{Q}$-divisor Δ on X is called effective if there is a positive integer k such that kD is a divisor and $\mathscr{O}_X(kD)$ has a non-zero global section.

A fibration $p : X \to Z$ between normal quasi-projective varieties is a dominant proper morphism such that every fiber is connected, that is $p_*\mathscr{O}_X \cong \mathscr{O}_Z$. If $X \to Z$ is just a proper morphism, then we have the Stein factorisation $X \to V \to Z$ such that $X \to V$ is a fibration and $V \to Z$ is finite (see [Har77, Cor. III. 11.5]). A Fano fibration is a fibration $X \to Z$ such that $-K_X$ is relatively ample. A Mori fibration is a Fano fibration such that the relative Picard number is 1.

Let $p : X \to Z$ be a morphism and D be a prime $\mathbb{Q}$-Cartier Weil divisor in Z. Let k be a positive integer such that kD is Cartier. Denote the $\mathbb{Q}$ divisor $\frac{1}{k}p^*(kD)$ by p^*D. We define the multiplicity $m(p, D) =\min\{\text{coefficient of } E \text{ in } p^*D \mid E \text{ is an irreducible component of } p^*D \text{ which dominates } D\}$. There are only finitely many divisors D such that $m(p, D) > 1$. Moreover, if $p : X \to Z$ is a morphism from a normal variety to a smooth curve, then we define the ramification divisor by $R = \sum_{z\in Z} p^*z - (p^*z)_{red}$ where $(p^*z)_{red}$ is the sum of the irreducible components of p^*z.

2.1. **Reflexive sheaves on normal varieties.** In this subsection, we gather some properties of reflexive sheaves. For a coherent sheaf $\mathscr{F}$ on a normal variety X, we denote by $\mathscr{F}^{**}$ the double dual of $\mathscr{F}$. The sheaf $\mathscr{F}$ is reflexive if and only if $\mathscr{F} \cong \mathscr{F}^{**}$. In particular, $\mathscr{F}^{**}$ is reflexive and we also call it the reflexive hull of $\mathscr{F}$. Denote $(\mathscr{F}^{\otimes m})^{**}$ by $\mathscr{F}^{[\otimes m]}$ and $(\bigwedge^m \mathscr{F})^{**}$ by $\mathscr{F}^{[\wedge m]}$ for any $m > 0$. For two coherent sheaves $\mathscr{F}$ and $\mathscr{G}$, let $\mathscr{F}[\otimes]\mathscr{G} = (\mathscr{F} \otimes \mathscr{G})^{**}$. The following proposition is an important criterion for reflexive sheaves on normal varieties.

Proposition 2.1 ([Har80, Prop. 1.6])**.** *Let $\mathscr{F}$ be a coherent sheaf on a normal variety X. Then $\mathscr{F}$ is reflexive if and only if $\mathscr{F}$ is torsion-free and for each open $U \subseteq X$ and each closed subset $Y \subseteq U$ of codimension at least 2, $\mathscr{F}|_U \cong j_*(\mathscr{F}|_{U\setminus Y})$, where $j : U \setminus Y \to U$ is the inclusion map.*

As a corollary of this proposition, we prove the following lemma.

Lemma 2.2. *If $\varphi : X \dashrightarrow X'$ is a birational map between normal projective varieties such that φ^{-1} does not contract any divisor, then we have a natural injection $H^0(X, (\Omega^1_X)^{[\otimes m]}) \hookrightarrow H^0(X', (\Omega^1_{X'})^{[\otimes m]})$ for any integer $m > 0$.*

Proof. Since the birational map $\varphi^{-1} : X' \dashrightarrow X$ does not contract any divisor, it induces an isomorphism from an open subset W' of X' such that $\mathrm{codim} X'\setminus W \geqslant 2$ onto an open subset W of X. By Proposition 2.1, we have $H^0(X', (\Omega^1_{X'})^{[\otimes m]}) \cong H^0(W', (\Omega^1_{W'})^{[\otimes m]})$ for any $m > 0$. Moreover, since W is an open subset of X, we obtain $H^0(X, (\Omega^1_X)^{[\otimes m]}) \hookrightarrow H^0(W, (\Omega^1_W)^{[\otimes m]}) \cong H^0(W', (\Omega^1_{W'})^{[\otimes m]}) \cong H^0(X', (\Omega^1_{X'})^{[\otimes m]})$ for all $m > 0$. □

The proof of the following lemma is left to the reader.

Lemma 2.3. *Let $0 \to \mathscr{A} \to \mathscr{B} \to \mathscr{C} \to 0$ be an exact sequence of locally free sheaves on a variety X. Then for any $m > 0$, we have a filtration of locally free sheaves $\mathscr{B}^{\otimes m} = \mathscr{R}_0 \supseteq \cdots \supseteq \mathscr{R}_{m+1} = 0$ such that $\mathscr{R}_i/\mathscr{R}_{i+1}$ is isomorphic to the direct sum of copies of $\mathscr{A}^{\otimes i} \otimes \mathscr{C}^{\otimes m-i}$ for all $0 \leq i \leq m$.*

From this lemma, we can deduce the following lemma which is very important in the paper.

Lemma 2.4. *Let $\mathscr{A} \to \mathscr{B} \to \mathscr{C}$ be a complex of coherent sheaves on a normal variety X. Assume that there is an open subset W of X with $\mathrm{codim} X\setminus W \geqslant 2$ such that we have an exact sequence of locally free sheaves $0 \to \mathscr{A}|_W \to \mathscr{B}|_W \to \mathscr{C}|_W \to 0$ on W. If $H^0(X, \mathscr{A}^{\otimes r}[\otimes]\mathscr{C}^{\otimes t}) = \{0\}$ for all $t > 0$ and $r \geqslant 0$, then $H^0(X, \mathscr{B}^{[\otimes m]}) \cong H^0(X, \mathscr{A}^{[\otimes m]})$ for all $m > 0$.*

Proof. On W, we have $H^0(W, \mathscr{A}|^{\otimes r}_W \otimes \mathscr{C}|^{\otimes t}_W) = \{0\}$ for all $t > 0$ and $r \geqslant 0$ by Proposition 2.1. Fix $m > 0$, then we have a filtration $\mathscr{B}|^{\otimes m}_W = \mathscr{R}_0 \supseteq \cdots \supseteq \mathscr{R}_{m+1} = 0$ such that $\mathscr{R}_i/\mathscr{R}_{i+1}$ is isomorphic to the direct sum of copies of $\mathscr{A}|^{\otimes i}_W \otimes \mathscr{C}|^{\otimes m-i}_W$ for all $0 \leq i \leq m$. Since $H^0(X, \mathscr{A}|^{\otimes r}_W \otimes \mathscr{C}|^{\otimes t}_W) = \{0\}$ for all $t > 0$ and $r \geqslant 0$, we have $H^0(W, \mathscr{R}_i) \cong H^0(W, \mathscr{R}_{i+1})$ for $0 \leqslant i \leqslant m - 1$. Thus $H^0(W, \mathscr{B}|^{\otimes m}_W/\mathscr{A}|^{\otimes m}_W) = \{0\}$ and $H^0(W, B|^{\otimes m}_W) \cong H^0(W, A|^{\otimes m}_W)$. By Proposition 2.1, we have $H^0(X, \mathscr{B}^{[\otimes m]}) \cong H^0(X, \mathscr{A}^{[\otimes m]})$. □

One of the applications of the lemma above is the following, which gives a relation between pluri-forms and fibrations.

Lemma 2.5. *Let $p : X \to Z$ be an equidimensional morphism between normal varieties. Assume that general fibers of p do not carry any non-zero pluri-form. Then $H^0(X, (\Omega^1_X)^{[\otimes m]}) \cong H^0(X, ((p^*\Omega^1_Z)^{sat})^{[\otimes m]})$ for $m > 0$, where $(p^*\Omega^1_Z)^{sat}$ is the saturation of the image of $p^*\Omega^1_Z$ in $\Omega^{[1]}_X$.*

Proof. We have an exact sequence of coherent sheaves $0 \to \mathscr{F} \to \Omega_X^{[1]} \to \mathscr{G} \to 0$ on X, where $\mathscr{F} = (p^*\Omega_Z^1)^{sat}$ and $\mathscr{G}$ is a torsion free sheaf. Let V be the smooth locus of Z and let $W = p^{-1}(V)$. If W_{ns} is the largest open subset of W on which $\mathscr{F}$, Ω_X^1 and $\mathscr{G}$ are locally free, then $\mathrm{codim} X\backslash W_{ns} \geqslant 2$ and we have an exact sequence $0 \to \mathscr{F}|_{W_{ns}} \to \Omega^1_{W_{ns}} \to \mathscr{G}|_{W_{ns}} \to 0$ of locally free sheaves on W_{ns}.

If F is a general fiber of $p|_{W_{ns}}$, then $\mathscr{F}|_F$ is the direct sum of $\mathscr{O}_F$ and $\mathscr{G}|_F$ is isomorphic to Ω_F^1. Since general fibers of p do not carry any non-zero pluri-form, neither does F by Proposition 2.1. Hence we have $H^0(W_{ns}, \mathscr{F}|_{W_{ns}}^{\otimes r} \otimes \mathscr{G}|_{W_{ns}}^{\otimes t}) = \{0\}$ for all $t > 0$ and $r \geqslant 0$. By Lemma 2.4, we have $H^0(W_{ns}, \mathscr{F}|_{W_{ns}}^{\otimes m}) \cong H^0(W_{ns}, (\Omega^1_{W_{ns}})^{\otimes m})$ for all $m > 0$. Since $\mathrm{codim} X\backslash W_{ns} \geqslant 2$, by Proposition 2.1, we have $H^0(X, (\Omega_X^1)^{[\otimes m]}) \cong H^0(X, \mathscr{F}^{[\otimes m]})$ for all $m > 0$. □

If X is a normal variety and E is a Weil divisor on X, then E is a Cartier divisor on the smooth locus X_{ns} of X and it induces an invertible sheaf $\mathscr{O}_{X_{ns}}(E)$ on X_{ns}. Let $\mathscr{O}_X(E)$ be the push-forward of $\mathscr{O}_{X_{ns}}(E)$ on X. Then by Proposition 2.1, $\mathscr{O}_X(E)$ is a reflexive sheaf on X since X is smooth in codimension 1. Conversely, if $\mathscr{F}$ is a reflexive sheaf of rank 1 on X, it is an invertible sheaf in codimension 1. There is a Weil divisor D on X such that $\mathscr{F} \cong \mathscr{O}_X(D)$. If X is $\mathbb{Q}$-factorial, then for any 1-cycle α on X, we can define the intersection number $\mathscr{F} \cdot \alpha = D \cdot \alpha = \frac{1}{k}(kD) \cdot \alpha$, where k is a positive integer such that kD is Cartier. This expression is independent to the choice of D (see [Rei79, Appendix to §1] for more details).

2.2. **Minimal model program.** We will recall some basic definition and properties of the minimal model program (MMP for short). A pair (X, Δ) consists of a normal quasi-projective variety X and a boundary Δ, *i.e.* a $\mathbb{Q}$-Weil divisor divisor $\Delta = \Sigma_{j=1}^k d_j D_j$ on X such that the D_j's are pairwise distinct prime divisors and all d_j are contained in $[0, 1]$. Recall the definition of singularities of pairs.

Definition 2.6 ([KM98, Def. 2.34])**.** Let (X, Δ) be a pair with $\Delta = \Sigma_{j=1}^k d_j D_j$. Let $r : \widetilde{X} \to X$ be a log resolution of singularities of (X, Δ). Assume that $K_X + \Delta$ is $\mathbb{Q}$-Cartier, then we can write $K_{\widetilde{X}} + r_*^{-1}\Delta = r^*(K_X + \Delta) + \Sigma a_i E_i$, where E_i are r-exceptional divisors. We call a_i the discrepancy of E_i with respect to (X, Δ). Then (X, Δ) is terminal (resp. canonical, klt) if $a_i > 0$ (resp. $a_i \geqslant 0$, $a_i > -1$ and $d_j < 1$ for all j) for all i.

Remark 2.7. If X is terminal, then it is smooth in codimension 2. If X is canonical, then K_X is Cartier in codimension 2 (see [KM98, Cor. 5.18]).

For a klt pair (X, Δ) such that X is $\mathbb{Q}$-factorial, we can run a $(K_X + \Delta)$-MMP and obtain a sequence of rational maps $X = X_0 \dashrightarrow X_1 \dashrightarrow \cdots$ such that (X_i, Δ_i) is a klt pair and X_i is $\mathbb{Q}$-factorial, where Δ_i is the strict transform of Δ. Every elementary step in a MMP is either a divisorial contraction which is a morphism contracting an irreducible divisor or a flip which is an isomorphism in codimension 1. More generally, if $h : X \to T$ is a morphism, we can run a h-relative MMP such that for all i, there is a morphism $h_i : X_i \to T$ with $h_i \circ \varphi_i = h$, where φ_i is the birational map $X \dashrightarrow X_i$. For more details on MMP, we refer to [KM98, §3]. One of the most important problems in MMP is that if the sequence of rational maps above terminates. If it does, we will get the result of this h-relative MMP $X \dashrightarrow X'$. Let Δ' be the direct image of Δ. Then either $K_{X'} + \Delta'$ is relative nef over T or we have a $(K_{X'} + \Delta')$-Mori fibration $g : X' \to Z$ over T such that $-(K_{X'} + \Delta')$ is g-ample. Thanks to [Kaw92, Thm. 1], we know that any MMP for a klt pair (X, Δ) such that $\dim X \leqslant 3$ terminates.

Lemma 2.8. *Let $\varphi : X \dashrightarrow X'$ be an extremal divisorial contraction or a flip such that both X and X' are projective. Assume that X has $\mathbb{Q}$-factorial canonical singularities. If Y is an irreducible closed subvariety in X' such that it is the center of a divisor E over X' which has discrepancy 0, then φ^{-1} is a morphism around the generic point of Y.*

Proof. Assume the opposite. Then the discrepancy of E in X is strictly smaller than the one in X' (see [KM98, Lem. 3.38]). This implies that X does not have canonical singularities along the center of E in X which is a contradiction. □

Proposition 2.9. *Let X be a projective threefold which has at most canonical singularities. Let X' be the result of a MMP for X and denote the birational map $X \dashrightarrow X'$ by f_X. Let Γ be the normalisation of the graph of f_X. Then there is a natural isomorphism $H^0(X, (\Omega_X^1)^{[\otimes m]}) \cong H^0(\Gamma, (\Omega_\Gamma^1)^{[\otimes m]})$ for all $m > 0$.*

Proof. Since there is a natural birational projection $pr_1 : \Gamma \to X$, we have an injection $H^0(\Gamma, (\Omega^1_\Gamma)^{[\otimes m]}) \hookrightarrow H^0(X, (\Omega^1_X)^{[\otimes m]})$ by Lemma 2.2. Let $pr_2 : \Gamma \to X'$ be the natural projection. Now let $\sigma_X \in H^0(X, (\Omega^1_X)^{[\otimes m]})$ be a non-zero element. Since X, Γ and X' are birational, σ_X induces a rational section σ_Γ of $(\Omega^1_\Gamma)^{[\otimes m]}$ and an element $\sigma_{X'}$ of $H^0(X', (\Omega^1_{X'})^{[\otimes m]})$ (see Lemma 2.2). In order to prove that σ_Γ is a regular section it is enough to prove that σ_Γ does not have pole along any pr_1-exceptional divisor. Let E be an exceptional divisor for pr_1. Let $C \subseteq E$ be a curve which is exceptional for pr_1, then C is not contracted by pr_2 since the graph of f_X is included in $X \times X'$ and the normalisation map is finite. Hence $Y = pr_2(E) \subseteq X'$ is a curve and f_X^{-1} is not regular around the generic point of Y. By Lemma 2.8, X' has terminal singularities around the generic point of Y. Hence it is smooth around the generic point of Y since $\mathrm{codim}_{X'} Y = 2$ (see [KM98, Cor. 5.18]). Hence σ_Γ does not have pole along E (σ_Γ is the pullback of $\sigma_{X'}$ by pr_2). □

2.3. **Examples.** In this subsection, we will give some examples of rationally connected varieties which carry non-zero pluri-forms. First we will give an example of such varieties with terminal singularities. For the construction of this example, we use the method of Theorem 3.2.

Example 2.10. Let $C_1 = \{[a : b : c] \in \mathbb{P}^2 \mid a^3 + (a+c)b^2 + c^3 = 0\}$ be a smooth elliptic curve in $\mathbb{P}^2$. Let $X_1 = \{([a : b : c], [x : y : z : t]) \in \mathbb{P}^2 \times \mathbb{P}^3 \mid a^3 + (a+c)b^2 + c^3 = 0,\ (a^2 + 2b^2 + c^2)x^2 + (a^2 + 3b^2 + c^2)y^2 + (a^2 + 4b^2 + 3c^2)z^2 + (a^2 + 5b^2 + 6c^2)t^2 = 0\}$. Then X_1 is a smooth threefold and we have an induced fibration $\pi_1 : X_1 \to C_1$ such that general fibers of π_1 are smooth quadric surfaces which are Fano surfaces. Moreover, X_1 has Picard number 2 by the Lefschetz theorem (see [Laz04, Exmp. 3.1.23]). Hence π_1 is a Mori fibration. Moreover, since π_1 is smooth, we have $H^0(X_1, (\Omega^1_{X_1})^{\otimes 2}) \cong H^0(C_1, (\Omega^1_{C_1})^{\otimes 2})$ by Lemma 2.5. Let G be the group $\mathbb{Z}/2\mathbb{Z}$ and let $g \in G$ be the generator. We have an action of G on $\mathbb{P}^2 \times \mathbb{P}^3$ defined by $g \cdot ([a : b : c], [x : y : z : t]) = ([a : -b : c], [-x : -y : z : t])$. This action induces an action of G on X_1 and an action of G on C_1 such that π_1 in G-equivariant. We have $C_1/G = \mathbb{P}^1$. The action of G on X_1 is étale in codimension 2 and it has exactly 16 fix points. Thus $X = X_1/G$ is a threefold which has at most $\mathbb{Q}$-factorial terminal singularities by [Rei87, (5.1)]. We have a fibration $\pi : X \to \mathbb{P}^1$ induced by π_1. Moreover, the $\mathbb{Q}$-dimension of $\mathrm{Pic}(X)_\mathbb{Q}$ is not larger than the Picard number of X_1. Hence π is a Mori fibration since general fibers of π are smooth quadric surfaces. Hence X is rationally connected by [GHS03, Thm. 1.1]. In addition, $H^0(X, (\Omega^1_X)^{[\otimes 2]}) \cong H^0(X_1, (\Omega^1_{X_1})^{\otimes 2})^G$ since $X_1 \to X$ is étale in codimension 1. Moreover $H^0(C_1, (\Omega^1_{C_1})^{\otimes 2})^G \cong H^0(\mathbb{P}^1, \mathscr{O}_{\mathbb{P}^1}) \cong \mathbb{C}$ by [Ou13, Prop. 6.2]. Hence we have $H^0(X, (\Omega^1_X)^{[\otimes 2]}) \cong \mathbb{C}$.

Remark 2.11. If X is a normal rationally connected threefold and if X' is the result of a MMP for X, then $H^0(X, (\Omega^1_X)^{[\otimes m]})$ may be strictly included in $H^0(X', (\Omega^1_{X'})^{[\otimes m]})$ for some $m > 0$. For example, let $X' \to \mathbb{P}^1$ be the variety described in the example above. Then X' carries non-zero pluri-forms. Let $X \to X'$ be a resolution of singularities of X'. Then X' is the result of a MMP for X since X' has terminal singularities. However, since X is rationally connected and smooth, we have $H^0(X, (\Omega^1_X)^{[\otimes m]}) = \{0\}$ for any $m > 0$ (see [Kol96, Cor. IV.3.8]).

We will give two examples of rationally connected threefolds with canonical singularities which carry non-zero pluri-forms. In both cases, there is always a Mori fibration from the threefold X to a surface Z. In Example 2.12, the base Z is a Calabi-Yau surface with klt singularities which carries non-zero pluri-forms. In Example 2.14, the base Z is isomorphic to $\mathbb{P}^2$ which does not carry any non-zero pluri-form.

Example 2.12. Let C be the curve in $\mathbb{P}^2$ defined by $C = \{[x_1 : x_2 : x_3] \in \mathbb{P}^2 \mid x_1^3 + x_2^3 + x_3^3 = 0\}$. There exists an action of group $G = \mathbb{Z}/3\mathbb{Z}$ on C defined by $g \cdot [x_1 : x_2 : x_3] = [\xi x_1 : x_2 : x_3]$ where g is a generator of G and ξ is a primitive 3rd root of unity. Let $Z_1 = C \times C$. Then there is an action of G on Z_1 induced by the one of G on C. We have $Z = Z_1/G$ is a klt rationally connected surface such that K_Z is a torsion (see [Tot10, Example 10]). In particular, K_Z is an effective $\mathbb{Q}$-Cartier divisor. Let $X_1 = \mathbb{P}^1 \times Z_1$ and define an action of G on $\mathbb{P}^1$ by $g \cdot [y_1 : y_2] \to [\xi y_1 : y_2]$ where $[y_1 : y_2]$ are homogeneous coordinates of $\mathbb{P}^1$. Then we have an action of G on the smooth threefold X_1 and $X = X_1/G$ has canonical singularities (see [Rei79, Thm. 3.1]). We also have an Mori fibration $X \to Z$ whose general fibers are isomorphic to $\mathbb{P}^1$. Since Z is rationally connected, so is X by [GHS03, Thm. 1.1]. Moreover, since K_Z is effective, we have $H^0(X, (\Omega^1_X)^{[\otimes m]}) \neq \{0\}$ for some $m > 0$ (see §4.2).

In the example above, the non-zero pluri-forms come from K_Z, the canonical divisor of the base surface. In the following example, $-K_Z$ is ample. However X still carries non-zero pluri-forms. These forms come

from the multiple fibers of the fibration $X \to Z$. Before giving the example, we will first introduce a method to construct fibers with high multiplicities.

Construction 2.13. We want to construct a fibration p from a normal threefold T with canonical singularities to a smooth surface S such that $p^*C = 2(p^*C)_{red}$, where C is a smooth curve in S.

Let $T_0 = \mathbb{P}^1 \times S$ where S is a smooth projective surface. Let pr_1, pr_2 be the natural projections $T_0 \to \mathbb{P}^1$ and $T_0 \to S$. Let $C \subseteq S$ be a smooth curve and let $C_0 = \{z\} \times C$ be a section of pr_2 over C in T_0, where z is a point of $\mathbb{P}^1$. Let $E_0 = pr_2^*C \subseteq T_0$ be a smooth divisor. We will perform a sequence of birational transformation of T_0.

First we blow up C_0, we obtain a morphism $T_1 \to T_0$ with exceptional divisor E_1. Denote still by E_0 the strict transform of E_0 in T_1. If F is a fiber of $T_1 \to S$ over a point of C, then $F = F_1 + F_0$ where $F_i \subseteq E_i$ are rational curves and $K_{T_1} \cdot F_i = -1$, $E_i \cdot F_i = -1$.

Now we blow up C_1, the intersection of E_0 and E_1. We obtain a morphism $T_2 \to T_1$ with exceptional divisor E_2. Denote still by E_0, E_1 the strict transforms of E_0 and E_1 in T_2. If F is any set-theoretic fiber of $T_2 \to S$ over a point of C, then $F = F_0 + F_1 + F_2$ where $F_i \subseteq E_i$ are rational curves and $K_{T_2} \cdot F_i = 0$, $E_i \cdot F_i = -2$ for $i = 0, 1$, $K_{T_2} \cdot F_2 = -1$, $E_2 \cdot F_2 = -1$. Let p_2 be the fibration $T_2 \to S$.

We will blow down E_1 and E_0 in T_2. Let H be an ample $\mathbb{Q}$-divisor such that $(H + E_0) \cdot F_0 = 0$. Since $E_0 \cdot F_1 = 0$, there is a positive rational number b such that $(H + E_0 + bE_1) \cdot F_1 = 0$. Moreover $(H + E_0 + bE_1) \cdot F_0 = 0$ since $E_1 \cdot F_0 = 0$. Let A be a sufficiently ample divisor on S. Then $D = p_2^*A + H + E_0 + bE_1$ is a nef and big $\mathbb{Q}$-divisor. Moreover, any curve B have intersection number 0 with D must be contract by $T_2 \to S$ since A is sufficiently ample. The curve B is also contained in $E_0 \cup E_1$ since H is ample. Since $K_{T_2} \cdot F_0 = K_{T_2} \cdot F_1 = 0$, there exists a large integer k such that $kD - K_{T_2}$ is nef and big. Then by the basepoint-free theorem (see [KM98, Thm. 3.3]), there is a large enough integer a such that aD is Cartier and $|aD|$ is basepoint-free. The linear system $|aD|$ induces a contraction which contracts E_0 and E_1.

By contracting E_0 and E_1, we obtain a threefold T. And there is an induced fibration $p : T \to S$ such that $p^*C = 2(p^*C)_{red}$. Notice that $T_2 \to T$ is a resolution of singularities and $K_{T_2} = f^*K_T$. Hence V has canonical singularities.

Now we will construct the example.

Example 2.14. Let $C = \{[x : y : z] \in \mathbb{P}^2 \mid x^6 + y^6 + z^6 = 0\}$ be a smooth curve in $\mathbb{P}^2$. Then $\mathscr{O}_{\mathbb{P}^2}(2K_{\mathbb{P}^2} + C) \cong \mathscr{O}_{\mathbb{P}^2}$ is effective. Let $X_0 = \mathbb{P}^1 \times \mathbb{P}^2$. By the method of Construction 2.13, we can construct a fiber space $p : X \to \mathbb{P}^2$ such that $p^*C = 2(p^*C)_{red}$. Then X has canonical singularities. Since general fibers of p are isomorphic to $\mathbb{P}^1$, X is rationally connected by [GHS03, Thm. 1.1]. Moreover, since $2(K_{\mathbb{P}^2} + \frac{1}{2}D)$ is an effective divisor and p is equidimensional, we have $H^0(X, (\Omega^1_X)^{[\otimes 4]}) \neq \{0\}$ (see §4.2).

Note that in the three examples above, the variety X we constructed is a rationally connected varieties with canonical singularities. The divisor K_X is not pseudo-effective and we always have a Mori fibration from X to a variety Z. The non-zero pluri-forms of X come from the base Z. However, in the following example, the variety we construct is a rationally connected threefold with klt singularities whose canonical divisor is ample. Its non-zero pluri-forms come from its canonical divisor.

Example 2.15. Let X_1 be the Fermat hypersurface in $\mathbb{P}^4$ defined $x_0^6 + x_1^6 + \cdots + x_4^6 = 0$. Then X_1 is a smooth threefold such that K_{X_1} is ample. Moreover, the Picard number of X_1 is 1 by the Lefschetz theorem (see [Laz04, Exmp. 3.1.23]). Let G be the group $\mathbb{Z}/6\mathbb{Z}$ with generator g. Define an action of G on $\mathbb{P}^4$ by $g \cdot [x_0 : \cdots : x_4] = [\omega x_0 : \omega x_1 : x_2 : x_3 : x_4]$, where ω is a primitive 6th root of unity. This action induces an action of G on X_1 which is free in codimension 1. Denote the quotient X_1/G by X and the natural morphism $X_1 \to X$ by π. Then X has $\mathbb{Q}$-factorial klt singularities but does not have canonical singularities (see [Rei79, Thm 3.1]). Now we will prove that X is rationally connected. First we will prove X is uniruled. If w is a general point in X, then there is a point $w_1 \in X_1$ such that $\pi(w_1) = w$ and the first two coordinates of w_1 are all non-zero. There are two hyperplanes in $\mathbb{P}^4$ passing through w_1, $H_1 = \{a_2x_2 + a_3x_3 + a_4x_4 = 0\}$ and $H_2 = \{b_2x_2 + b_3x_3 + b_4x_4 = 0\}$, such that the intersection $C_1 = H_1 \cap H_2 \cap X_1$ is a smooth curve of genus 10. Moreover, there are exactly 6 points on C_1, given by $X_1 \cap \{x_2 = x_3 = x_4 = 0\}$, which are fixed under the action of G. Hence $\pi|_{C_1} : C_1 \to \pi(C_1) = C$ is ramified exactly at those 6 points with degree 6. By adjunction formula, we have the genus of C, $g(C) = 1 + 2^{-1} \times 6^{-1} \times (2 \times 10 - 2 - 6 \times 5) = 0$. Hence C

is a smooth rational curve. This implies that X is uniruled. From the lemma below, we conclude that X is rationally connected.

Lemma 2.16. *Let X be a normal projective variety which is $\mathbb{Q}$-factorial. Assume that $\mathrm{Pic}_{\mathbb{Q}}(X) \cong \mathbb{Q}$. Then X is rationally connected if and only if it is uniruled.*

Proof. Note that if X is smooth, this was proven in [Cam92, Cor. 2.4]. Assume that X is uniruled and let $\pi : X \dashrightarrow Z$ be the MRC fibration for X (see [Cam92, Thm. 2.3]). We will argue by contradiction. Assume that $\dim X > \dim Z > 0$. By [Cam92, Thm. 2.3], there is a Zariski open subset Z_0 of Z such that π is regular and proper over Z_0. Let C be a curve contained in some fiber of $\pi|_{Z_0}$. Let D be a non-zero effective Cartier divisor on Z_0. Let $E_0 = \pi^* D$. Then E_0 is a non-zero effective divisor on $X_0 = \pi^{-1} Z_0$. Let E be the closure of E_0 in X, then E is a non-zero effective Weil divisor on X. However, the intersection number of E and C is zero, which contradicts $\mathrm{Pic}_{\mathbb{Q}}(X) = \mathbb{Q}$. □

3. Fibrations over curves

3.1. **Construction of rationally connected varieties carrying non-zero pluri-forms.** In this subsection, we will give a method to construct a rationally connected varieties which carry non-zero pluri-forms (Theorem 3.2). Together with Theorem 1.4, we will see that every rationally connected threefold with $\mathbb{Q}$-factorial terminal singularities which carries non-zero pluri-forms can be constructed by this method.

From Lemma 2.5, we know that if general fibers of an equidimensional fibration $p : X \to Z$ do not carry any non-zero pluri-form, then $H^0(X, (\Omega^1_X)^{[\otimes m]}) \cong H^0(X, ((p^*\Omega^1_Z)^{sat})^{[\otimes m]})$ for all $m > 0$. Moreover, $H^0(X, ((p^*\Omega^1_Z)^{sat})^{[\otimes m]}) \cong H^0(Z, p_*((p^*\Omega^1_Z)^{sat})^{[\otimes m]}))$. Hence we would like to know what $p_*((p^*\Omega^1_Z)^{sat})^{[\otimes m]})$ is. In the case where Z is a smooth curve, this is not difficult to compute. Notice that if R is the ramification divisor of p, then $((p^*\Omega^1_Z)^{sat})^{[\otimes m]} \cong (p^*\Omega^1_Z)^{\otimes m} \otimes \mathscr{O}_X(mR)$ for $m > 0$. By the projection formula, we know that $p_*((p^*\Omega^1_Z)^{sat})^{[\otimes m]}) \cong (\Omega^1_Z)^{\otimes m} \otimes p_*\mathscr{O}_X(mR)$. Hence it is enough to compute $p_*\mathscr{O}_X(mR)$.

Lemma 3.1. *Let $p : X \to Z$ be a fibration from a normal variety to a smooth curve. Let z be a point in Z. Let $D = p^*z - (p^*z)_{red}$. Then $p_*\mathscr{O}_X(mD) \cong \mathscr{O}_Z([\frac{m(m(p,z)-1)}{m(p,z)}]z)$ for any $m \geqslant 0$.*

Proof. The problem is local and we may assume that Z is affine. We know that $p_*\mathscr{O}_W(mD)$ is an invertible sheaf. If θ is a section of $p_*\mathscr{O}_W(mD)$, then it is a rational function on Z whose pullback on W is a rational function which can only have pole alone p^*z. Hence θ can only have pole at z. Let d_θ be the degree of the pole, then the pullback of θ in V is a section of $\mathscr{O}_W(mD)$ if and only if $m(p,z)d_\theta \leqslant m(m(p,z)-1)$, *i.e.* $d_\theta \leqslant [\frac{m(m(p,z)-1)}{m(p,z)}]$. □

This relation between ramification divisor and pluri-forms gives us an idea of how to construct rationally connected varieties which carry non-zero pluri-forms. Notice that if $p : X \to \mathbb{P}^1$ is a fibration such that general fibers of p are rationally connected, then X is rationally connected by [GHS03, Thm. 1.1]. Moreover, if general fibers of p do not carry any non-zero pluri-form, then from the discussion above X carries non-zero pluri-forms if and only if $(\Omega^1_{\mathbb{P}^1})^{\otimes m} \otimes p_*\mathscr{O}_X(mR)$ has non-zero sections for some $m > 0$, where R is the ramified divisor of p. However, $\Omega^1_{\mathbb{P}^1} \cong \mathscr{O}_{\mathbb{P}^1}(-2)$ and $p_*\mathscr{O}_X(mR) \cong \mathscr{O}_{p^1}(\sum_{z\in\mathbb{P}^1}[\frac{(m(p,z)-1)m}{m(p,z)}])$ since any two points in $\mathbb{P}^1$ are linearly equivalent. Hence X carries non-zero pluri-forms if and only if $\sum_{z\in\mathbb{P}^1} \frac{m(p,z)-1}{m(p,z)} \geqslant 2$.

Now we will try to construct this kind of varieties by taking quotients. Let T be a normal projective variety, let B be a smooth projective curve and let G be a finite commutative group. Assume that there are actions of G on T and B. Assume that there is a G-equivariant fibration $q : T \to B$. Let $p_T : T \to T/G$, $p_B : B \to B/G$ be natural projections and let $h : T/G \to B/G$ be the induced fibration. Let S_i, $i = 1, ..., r$ be all the G-orbits in B whose cardinality is less than $|G|$, the cardinality of G. Let z_i be the image of S_i under the map $B \to B/G$. Then the z_i's are the points in B/G over which $B \to B/G$ is ramified. Let G_i be the stabilizer of a point b_i in S_i. Then G_i acts on the set-theoretic fiber $q^{-1}\{b_i\} = F_i$. If A_i is a component in F_i and the stabilizer of a general point in A_i has cardinality d_i, then p_T is ramified along $G_i \cdot A_i$ of degree d_i, where $G_i \cdot A_i$ is the orbit of A_i under G_i. In this case, $p_W(A_i)$ has coefficient $\frac{e_i f_i}{d_i}$ in h^*z_i, where e_i is the coefficient of A_i in q^*b_i and f_i is the cardinality of G_i. Denote $\min\{\frac{e_i f_i}{d_i} \mid A_i \text{ a component in } F_i\}$ by s_i. Then s_i is equal to $m(h, z_i)$. We have the theorem below.

Theorem 3.2. *Let T be a projective normal variety, B be a projective smooth curve with positive genus and G be a finite commutative group. Assume they satisfy the following:*

(1) *There is a fibration q from T to B such that for any general fiber F_q of q, F_q is rationally connected and does not carry any non-zero pluri-form. Moreover, $m(q,b)=1$ for all $b\in B$.*
(2) *There exist actions of G on B and on T such that q is G-equivariant.*
(3) *The quotient B/G is isomorphic to $\mathbb{P}^1$.*

Then the quotient $X=T/G$ is a normal projective rationally connected variety and there is a fibration $p:X\to\mathbb{P}^1$. Moreover, if we define s_i, $i=1,...,r$ as above, then X carries non-zero pluri-forms if and only if $\Sigma_{i=1}^r\frac{s_i-1}{s_i}\geqslant 2$. In fact, $H^0(X,(\Omega^1_X)^{[\otimes m]})\cong H^0(\mathbb{P}^1,\mathscr{O}_{\mathbb{P}^1}(-2m+\sum_{i=1}^r[\frac{(s_i-1)m}{s_i}]))$ for $m>0$.

Proof. The fibration $p:X\to\mathbb{P}^1$ is induced by q. Since general fibers of q are rationally connected, general fibers of p are also rationally connected. Hence X is rationally connected by [GHS03, Thm. 1.1]. If F_p is a general fiber of p, then F_p does not carry any non-zero pluri-form neither. Hence, $H^0(X,(\Omega^1_X)^{[\otimes m]})\cong H^0(\mathbb{P}^1,\mathscr{O}_{\mathbb{P}^1}(-2m)\otimes p_*\mathscr{O}_X(mR))$ for $m>0$, where R is the ramification divisor. However, since $m(q,b)=1$ for all $b\in B$, we have $\sum_{i=1}^r[\frac{(s_i-1)m}{s_i}]=\sum_{z\in\mathbb{P}^1}[\frac{(m(p,z)-1)m}{m(p,z)}]$ by the definition of s_i. Finally, by Lemma 3.1, we obtain $p_*\mathscr{O}_X(mR)\cong\mathscr{O}_{\mathbb{P}^1}(\sum_{i=1}^r[\frac{(s_i-1)m}{s_i}])$ and $H^0(X,(\Omega^1_X)^{[\otimes m]})\cong H^0(\mathbb{P}^1,\mathscr{O}_{\mathbb{P}^1}(-2m+\sum_{i=1}^r[\frac{(s_i-1)m}{s_i}]))$. □

Remark 3.3. Conversely, Let $p:X\to\mathbb{P}^1$ be a fibration such that general fibers of p are rationally connected and do not carry non-zero pluri-form. If X carries non-zero pluri-forms, then X can be constructed by the method described above. In fact, by the discussion above, we have $\sum_{z\in\mathbb{P}^1}\frac{(m(p,z)-1)}{m(p,z)}\geqslant 2$. In particular, there are at least three points in $\mathbb{P}^1$ such that the multiplicity of p is larger than 1 over these points. Let $z_1,...,z_r$ be all the points in $\mathbb{P}^1$ over which the multiplicity of p is larger than 1. Let $m_i=m(p,z_i)$. Since $r\geqslant 3$, there is a smooth curve B and a Galois cover $p_B:B\to\mathbb{P}^1$ with Galois group G such that p_B is ramified exactly over the $z_i's$ and the degree of ramification is m_i at each point over z_i (see [KO82, Lem. 6.1]). Let T be the normalisation of the fiber product $X\times_{\mathbb{P}^1}B$. Then we obtain a natural fibration $q:T\to B$ such that $m(q,b)=1$ for all $b\in B$. Moreover, G acts naturally on T and $T/G\cong X$.

3.2. **The case of threefolds with canonical singularities.** If $p:X\to Z$ is a Mori fibration from a threefold which has $\mathbb{Q}$-factorial canonical singularities to a smooth curve, then general fibers of p are Fano surfaces which have canonical singularities. These surfaces do not carry any non-zero pluri-form by the following proposition.

Proposition 3.4. *If S is a Fano surface which has canonical singularities, then $H^0(S,(\Omega^1_S)^{[\otimes m]})=\{0\}$ for all $m>0$.*

Proof. Assume the opposite. If S' is the result of a MMP for S, then S' is a Mori fiber space. By [Ou13, Thm. 2.1], S' has Picard number 2 and we have a Mori fibration $S'\to\mathbb{P}^1$. Let p_S be the composition of $S\to S'\to\mathbb{P}^1$. By [Ou13, Thm1.4], there is a smooth curve B of positive genus and a finite morphism $B\to\mathbb{P}^1$ such that the natural morphism $S_B\to S$ is étale in codimension 1, where S_B is the normalisation of $S\times_{\mathbb{P}^1}B$. Hence S_B is a Fano surface. Moreover it has canonical singularities (see [KM98, Prop. 5.20]). So it is rationally connected by [HM07, Cor. 1.3 and 1.5]. Hence B is also rationally connected, which is a contradiction since its genus is positive. □

Thanks to this proposition, we obtain the following result.

Proposition 3.5. *Let X be a rationally connected threefold with $\mathbb{Q}$-factorial canonical singularities. Assume that X carries non-zero pluri-forms. Let X' be the result of a MMP for X and assume that there is a Mori fibration $p':X'\to\mathbb{P}^1$. Let Y be the normalisation of the graph of $X\dashrightarrow X'$. Then Y can be constructed by the method of Theorem 3.2. After a contraction from Y, we can obtain X.*

Proof. By Proposition 3.4 and Lemma 2.5, we know that every element in $H^0(X',(\Omega^1_{X'})^{[\otimes m]})$ comes from the base $\mathbb{P}^1$, so does every element in $H^0(X,(\Omega^1_X)^{[\otimes m]})$ by Lemma 2.2. But we only have a rational map $X\dashrightarrow X'\to\mathbb{P}^1$. However, by Proposition 2.9, we know that $H^0(X,(\Omega^1_X)^{[\otimes m]})\cong H^0(Y,(\Omega^1_Y)^{[\otimes m]})$. Moreover, we have a natural fibration $g:Y\to\mathbb{P}^1$ which is the composition of $Y\to X'\to\mathbb{P}^1$. If F_g is a general fiber of g, then there is a birational morphism $F_g\to S$, where S is a general fiber of $p':X'\to\mathbb{P}^1$. Since S does not carry any non-zero pluri-form by Proposition 3.4, neither does F_g by Lemma 2.2. From Remark 3.3, we know that Y can be constructed by the method of Theorem 3.2. After a contraction from Y, we obtain the variety X. □

4. Mori fibrations and non-zero pluri-form

In this section, we study relations between Mori fibrations and non-zero pluri-forms. We will consider Mori fibrations from a normal threefold to a normal surface. First, we recall the definition and some properties of slopes. Let X be a normal projective $\mathbb{Q}$-factorial variety. Let α be a class of 1-cycles in X. For a coherent sheaf $\mathscr{F}$ of positive rank, we define the slope $\mu_\alpha(\mathscr{F})$ of $\mathscr{F}$ with respect to α by

$$\mu_\alpha(\mathscr{F}) := \frac{\det(\mathscr{F})\cdot\alpha}{\mathrm{rank}(\mathscr{F})},$$

where $\det(\mathscr{F})$ is the reflexive hull of $\bigwedge^{\mathrm{rank}\mathscr{F}}\mathscr{F}$. A class of movable curves is a class of 1-cycles which has non negative intersection number with any pseudo-effective divisor. If α is a class of movable curves, then $\mu_\alpha^{max}(\mathscr{F}) = \sup\{\mu_\alpha(\mathscr{G}) \mid \mathscr{G}\subseteq\mathscr{F}$ a coherent subsheaf of positive rank$\}$ is well defined (see [MP97, p.42]). For any coherent sheaf $\mathscr{F}$, there is a saturated coherent subsheaf $\mathscr{G}\subseteq\mathscr{F}$ such that $\mu_\alpha^{max}(\mathscr{F}) = \mu_\alpha(\mathscr{G})$. If $\mathscr{E}$ and $\mathscr{F}$ are two coherent sheaves of positive rank, then $\mu_\alpha(\mathscr{F}\otimes\mathscr{E}) = \mu_\alpha(\mathscr{F}) + \mu_\alpha(\mathscr{E})$.

4.1. **General properties.** Consider a Mori fibration $p : X \to Z$ from a normal rationally connected threefold to a normal variety Z of positive dimension. Assume that X has $\mathbb{Q}$-factorial klt singularities. Let $D_1, ..., D_k$ be all prime Weil divisors in Z such that $m_i = m(p, D_i) > 1$. Let $\Delta = \sum_{i=1}^k \frac{m_i-1}{m_i}D_i$. Then $\det((p^*\Omega^1_Z)^{sat}) \cong \mathscr{O}_X(p^*(K_Z+\Delta))$ (see Remark 4.5). Assume that X carries non-zero pluri-forms and general fibers of p do not carry any non-zero pluri-form. If $\dim Z = 1$, then $K_Z+\Delta$ is an effective $\mathbb{Q}$-divisor of degree $-2+\sum_{i=1}^k \frac{m_i-1}{m_i}$ on $\mathbb{P}^1$ (see §3). The aim of this subsection is to prove something analogue in the case where $\dim Z = 2$. We prove that if $K_Z+\Delta$ is not pseudo-effective, then there will be a fibration form Z to $\mathbb{P}^1$. In this case, we have an induced fibration $X\to\mathbb{P}^1$ and we are in the same situation as in §3 (see Lemma 4.10). In order to do this, we will run a MMP for the pair (Z,Δ). To this end, we prove the following proposition which implies that the pair (Z,Δ) is klt.

Proposition 4.1. *Let $p : X \to Z$ be a Mori fibration from a $\mathbb{Q}$-factorial klt quasi-projective variety X to a normal variety Z. Let $D_1, ..., D_k$ be pairwise distinct prime Weil divisors in Z such that $p^*D_i = m_i(p^*D_i)_{red}$ with $m_i \geqslant 2$. Then the pair $(Z, \sum_{i=1}^k \frac{m_i-1}{m_i}D_i)$ is klt.*

Proof. Let $D = D_1 + \cdots + D_k$. By [KMM87, Lem. 5-1-5], Z is $\mathbb{Q}$-factorial. Notice that the problem is local in Z, we may assume that Z is affine. Let k_1 be the smallest positive integer such that k_1D_1 is a Cartier divisor. By taking the k_1-th root of the function defining the Cartier divisor k_1D_1, we can construct a finite covering $c_{1,1} : Z_{1,1} \to Z$ which is étale in codimension 1. Moreover, $c_{1,1}^*D_1$ is a prime Cartier divisor (see [Mor88, Prop.-Def. 1.11]). By taking the m_1-th root of the function defining the Cartier divisor $c_{1,1}^*D_1$, we can find a finite covering $c_{1,2} : Z_1 \to Z_{1,1}$ which is ramified exactly over D_1 with ramification degree m_1 (see [Laz04, §4.1B]). Let $c_1 = c_{1,2}\circ c_{1,1} : Z_1 \to Z$. Then $c_1^*D_1 = m_1(c_1^*D_1)_{red}$. Repeating this construction for the strict transform of D_2 in Z_1, we have a finite morphism $Z_2 \to Z_1$ which is ramified over the strict transform of D_2 in Z_1 with degree m_2. Let $c_2 : Z_2 \to Z$ be the composition of $Z_2 \to Z_1 \to Z$. Then c_2 is ramified over D_1+D_2 and $c_2^*D_i = m_i(c_1^*D_i)_{red}$ for $i = 1, 2$. By induction, we can construct a finite morphism $c_k : Z_k \to Z$ which is ramified over D such that $c_k^*D_i = m_i(c_1^*D_i)_{red}$ for all i. We have $K_{Z_k} = c_k^*(K_Z + \sum_{i=1}^k \frac{m_i-1}{m_i}D_i)$ is $\mathbb{Q}$-Cartier (for the proof of this formula, see Lemma 4.3 below). Let X_k be the normalisation of $X\times_Z Z_k$. Then the natural projection $c_X : X_k \to X$ is étale in codimension 1 and $K_{X_k} = c_X^*K_X$. Hence X_k is klt by [KM98, Prop. 5.20]. Moreover, $X_k \to Z_k$ is a Fano fibration since $X \to Z$ is a Mori fibration. Hence there is a $\mathbb{Q}$-divisor Δ_k such that the pair (Z_k, Δ_k) is klt by [Fuj99, Cor. 4.7]. Since K_{Z_k} is $\mathbb{Q}$-Cartier, Z_k is klt (see [KM98, Cor. 2.35]). So the pair $(Z, \sum_{i=1}^k \frac{m_i-1}{m_i}D_i)$ is also klt by [KM98, Prop. 5.20]. □

In the remaining of this subsection, our aim is to prove the following theorem.

Theorem 4.2. *Let $p : X \to Z$ be a Mori fibration from a projective normal threefold to a normal projective surface. Assume that X has $\mathbb{Q}$-factorial klt singularities. Let $D_1, ..., D_k$ be all prime Weil divisors in Z such that $m_i = m(p, D_i) > 1$. Let $\Delta = \sum_{i=1}^k \frac{m_i-1}{m_i}D_i$. If X carries non-zero pluri-forms and $K_Z+\Delta$ is not pseudo-effective, then the result Z' of any MMP for the pair (Z,Δ) has Picard number 2.*

First we would like to illustrate the idea of the proof in a simple case. Let $f_Z : (Z,\Delta) \to (Z',\Delta')$ be the result of a MMP for the pair (Z,Δ). Assume in a first stage that $Z = Z'$. We will argue by contradiction. Assume that Z' has Picard number 1. Since general fibers of p are isomorphic to $\mathbb{P}^1$, they don't carry any

non-zero pluri-form. By Lemma 2.5, the non-zero pluri-forms of X come from $(p^*\Omega^1_Z)^{sat}$. Since $K_Z+\Delta$ is not pseudo-effective, we can prove that there is a rank 1 reflexive subsheaf $\mathscr{H}$ of $(p^*\Omega^1_Z)^{sat}$ such that $\mathscr{H}^{[\otimes l]}$ is an invertible sheaf which has non-zero global sections for some positive integer l. Next, we can prove that $\mathscr{H}^{[\otimes l]}$ is isomorphic to $\mathscr{O}_X$ under the assumption that $\mathrm{Pic}_{\mathbb{Q}}Z=\mathbb{Q}$ (by using Lemma 4.8). Let $W\to X$ be the cyclic cover with respect to $\mathscr{H}$ and l. Let $W\to V\to Z$ be the Stein factorisation, then V will be a klt log Fano variety (by Lemma 4.7). Moreover $H^0(V,\Omega^{[1]}_V)\neq\{0\}$ (we use Lemma 4.9). This contradicts Theorem 1.2.

However, in general Z' is different from Z and the fibration $X\to Z'$ is not equidimensional. The idea of the proof will be the same as above but details will be more complicated. We will work over an open subset Z_0 of Z such that $f_Z|_{Z_0}$ is an isomorphism and $\mathrm{codim}\, Z'\backslash f_Z(Z_0)\geqslant 2$. For the complete proof of the theorem, we will need several lemmas.

Lemma 4.3. *Let $p:X\to Z$ be an equidimensional morphism between smooth varieties. Let n be the dimension of X and d be the dimension of Z. Let D be a prime divisor in Z and E be a prime divisor in X such that E is a component of p^*D. Assume that the coefficient of E in p^*D is k. Then for any general point $x\in E$, there is an open neighbourhood $U\subseteq X$ of x such that $(p^*\Omega^d_Z)^{sat}|_U\cong\mathscr{O}_X(p^*K_Z+(k-1)E)|_U$, where $(p^*\Omega^d_Z)^{sat}$ is the saturation of $p^*\Omega^d_Z$ in Ω^d_X.*

Proof. We may assume that E is smooth around x and D is smooth around $p(x)$. There exist local coordinates $(a_1,a_2,...,a_n)$ and $(b_1,b_2,...,b_d)$ of X and Z around x and $p(x)$ such that E is defined by $\{a_1=0\}$, D is defined by $\{b_1=0\}$ and p is given by $(a_1,a_2,...,a_n)\mapsto(a_1^k,a_2,...,a_d)$. With these coordinates, the natural morphism $p^*\Omega^1_Z\to\Omega^1_X$ is given by $(\mathrm{d}b_1,\mathrm{d}b_2,...,\mathrm{d}b_d)\mapsto(ka_1^{k-1}\mathrm{d}a_1,\mathrm{d}a_2,...,\mathrm{d}a_d)$ and the image of the natural morphism $p^*\Omega^d_Z\to\Omega^d_X$ is generated by $p^*(\mathrm{d}b_1\wedge\mathrm{d}b_2\wedge\cdots\wedge\mathrm{d}b_d)=(ka_1^{k-1}\mathrm{d}a_1)\wedge\mathrm{d}a_2\wedge\cdots\wedge\mathrm{d}a_d$. Hence $(p^*\Omega^d_Z)^{sat}$ is generated by $\mathrm{d}a_1\wedge\mathrm{d}a_2\wedge\cdots\wedge\mathrm{d}a_d$. Since $\{a_1=0\}$ defines the divisor E, we have $(p^*\Omega^d_Z)^{sat}|_U\cong\mathscr{O}_X(p^*K_Z+(k-1)E)|_U$ for some open neighbourhood U of x. □

Lemma 4.4. *Let $p:X\to Z$ be an equidimensional morphism between normal varieties. Assume that Z is $\mathbb{Q}$-factorial. Let $D_1,...,D_r$ be all the prime divisors in Z such that $m(p,D_i)$ is larger than 1. Write $\det((p^*(\Omega^1_Z))^{sat})\cong\mathscr{O}_X(M)$ where M is a divisor on X, then $M-(p^*(K_Z+\sum_{i=1}^r\frac{m(p,D_i)-1}{m(p,D_i)}D_i))$ is effective.*

Proof. By Proposition 2.1, we only have to prove the assertion on an open subset of X whose complement is of codimension at least 2. Hence we may assume that both X and Z are smooth and that $\sum_{i=1}^r D_i$ is smooth. In this case, by Lemma 4.3, the divisor M is linearly equivalent to $p^*K_Z+\sum_{i=1}^r\sum_{j=1}^{s_i}(n_{i,j}-1)E_{i,j}$ where the $E_{i,1},...,E_{i,s_i}$ are the components of p^*D_i and $n_{i,j}$ is the coefficient of $E_{i,j}$ in p^*D_i. Since $m(p,D_i)$ is the smallest integer among $n_{i,1},...,n_{i,s_i}$, we have $n_{i,j}-1\geqslant\frac{m(p,D_i)-1}{m(p,D_i)}\cdot n_{i,j}$. Thus $\sum_{i=1}^r\sum_{j=1}^{s_i}(n_{i,j}-1)E_{i,j}\geqslant\sum_{i=1}^r\frac{m(p,D_i)-1}{m(p,D_i)}p^*D_i$. □

Remark 4.5. With the notation above, if in addition p is a Mori fibration, then p^*D_i is irreducible for all i since p has relative Picard number 1. In this case we obtain that $\det((p^*(\Omega^1_Z))^{sat})$ is isomorphic to $\mathscr{O}_X(p^*(K_Z+\sum_{i=1}^r\frac{m(p,D_i)-1}{m(p,D_i)}D_i))$.

Lemma 4.6. *Let $p:X\to Z$ be an equidimensional fibration between normal varieties. Let $(p^*\Omega^1_Z)^{sat}$ be the saturation of the image of $p^*\Omega^1_Z$ in $\Omega^{[1]}_X$. Then $p_*(((p^*\Omega^1_Z)^{sat})^{[\wedge r]})\cong\Omega^{[r]}_Z$ for $r>0$.*

Proof. Fix $r>0$. Let $\mathscr{M}=p_*(((p^*\Omega^1_Z)^{sat})^{[\wedge r]})$. Let X_0 be the smooth locus of X and let $i:X_0\to X$ be the canonical injection. Then $\Omega^{[1]}_X=i_*\Omega^1_{X_0}$ by Proposition 2.1. Since $\mathrm{codim}X\backslash X_0\geqslant 2$ and Z is smooth in codimension 1, there is a smooth open subset Z_0 in Z which is contained in $p(X_0)$ and $\mathrm{codim}Z\backslash Z_0\geqslant 2$. By Proposition 2.1, it is enough to prove that $\mathscr{M}|_{Z_0}\cong\Omega^{[r]}_Z|_{Z_0}=\Omega^r_{Z_0}$. Hence, by replacing Z by Z_0, we may assume that Z is smooth and $p(X_0)=Z$. Let D be the sum of the prime divisors in Z whose pull-back by p are not reduced divisors. There is an open subset Z_1 of Z such that $D|_{Z_1}$ is snc and $\mathrm{codim}Z\backslash Z_1\geqslant 2$. From Proposition 2.1, by replacing Z by Z_2, we may assume that D is a snc divisor on Z.

First we will prove that there is a natural injection from Ω^r_Z to $\mathscr{M}$. In fact, on X_0, we have an injection from $p^*\Omega^1_Z|_{X_0}$ to $(p^*\Omega^1_Z)^{sat}|_{X_0}$. Hence there is an injection from $(p^*\Omega^1_Z)^{\wedge r}|_{X_0}$ to $((p^*\Omega^1_Z)^{sat})^{[\wedge r]}|_{X_0}$ on X_0. Since $(p^*\Omega^1_Z)^{\wedge r}$ is without torsion, we have an injection from $(p^*\Omega^1_Z)^{\wedge r}$ to $((p^*\Omega^1_Z)^{sat})^{[\wedge r]}$ on X. Since $p(X)=Z$, we obtain an injection from $p_*((p^*\Omega^1_Z)^{\wedge r})$ to $\mathscr{M}$. By the projection formula, this implies that $\Omega^r_Z\subseteq\mathscr{M}$.

Now we will prove that $\Omega^r_Z \cong \mathscr{M}$. Let $W = p^{-1}(Z\backslash D)$. Then the morphism $p|_W : W \to Z\backslash D$ is smooth in codimension 1. Thus, $((p^*\Omega^1_Z)^{sat})^{[\wedge r]}|_W \cong p^*\Omega^r_Z|_W$ (see the proof of Lemma 4.3). Then we obtain $\mathscr{M}|_{Z\backslash D} \cong \Omega^r_{Z\backslash D}$ by the projection formula. Let U be any open set in Z and let β be any element of $\mathscr{M}(U)$, *i.e.* a section of $\mathscr{M}|_U$. Then β is a rational section of $\Omega^r_Z|_U$ which can only have pole along D since $\mathscr{M}|_{Z\backslash D} \cong \Omega^r_{Z\backslash D}$. However, by the definition of $\mathscr{M}$, β induces a regular section of $\Omega^r_{X_0}|_{p^{-1}(U)}$. This implies that β does not have pole along D. Thus $\mathscr{M}(U) = \Omega^r_Z(U)$. Hence $\Omega^r_Z \cong \mathscr{M}$. □

Lemma 4.7. *Let $p : X \to Z$ be an equidimensional fibration from a normal variety X to a smooth variety Z. Let $D_1, ..., D_r$ be all the prime divisors in Z such that the multiplicity $m(p, D_i)$ is larger than 1. If $c_X : X_1 \to X$ is a finite morphism which is étale in codimension 1 and $X_1 \xrightarrow{p_1} Z_1 \xrightarrow{c_Z} Z$ is the Stein factorisation, then $K_{Z_1} \leqslant c_Z^*(K_Z + \sum_{i=1}^r \frac{m(p,D_i)-1}{m(p,D_i)} D_i)$. That is, there is an effective $\mathbb{Q}$-divisor Δ_1 on Z_1 such that $K_{Z_1} + \Delta_1 = c_Z^*(K_Z + \sum_{i=1}^r \frac{m(p,D_i)-1}{m(p,D_i)} D_i)$.*

Proof. If D is a prime divisor in Z, then $p_1^*c_Z^*D = c_X^*p^*D$. Let E be any irreducible component of c_Z^*D. Since c_X is étale in codimension 1, by comparing the coefficients of E in $p_1^*c_Z^*D = c_X^*p^*D$, the coefficient m_E of E in c_Z^*D is not larger than $m(p, D)$. This implies that $(m_E - 1) \leqslant (m(p, D) - 1)$. Now from Lemma 4.3, we obtain $K_{Z_1} = c_Z^*K_Z + \Sigma_{j=1}^s(m_j - 1)E_j$, where the $E_j's$ are all prime divisors along which c_Z^* is ramified and m_j is the degree of ramification of c_Z along E_j. By the discussion above, we have $(m_j - 1) \leqslant (m(p, p(E_j)) - 1) = m(p, p(E_j)) \cdot \frac{m(p,p(E_j))-1}{m(p,p(E_j))}$. Hence $K_{Z_1} \leqslant c_Z^*(K_Z + \sum_{i=1}^r \frac{m(p,D_i)-1}{m(p,D_i)} D_i)$. □

Lemma 4.8. *Let $p : X \to Z$ be a Mori fibration from a normal threefold X to a smooth surface Z. Let $D_1, ..., D_r$ be all the prime divisors in Z such that the multiplicity $m_i = m(p, D_i)$ is larger than 1. Assume that there is a projective $\mathbb{Q}$-factorial variety V such that we have an open embedding $j : Z \to V$ with $\mathrm{codim} V\backslash Z \geqslant 2$. Assume further that the pair $(V, \sum_{i=1}^r \frac{m_i-1}{m_i}\bar{D}_i)$ is klt, where $\bar{D}_i$ is the closure of $\bar{D}_i$ in V. If $\mathscr{F}$ is a rank 1 reflexive subsheaf of $\Omega^{[1]}_X$ such that $\mathscr{F}^{[\otimes m_0]} \cong \mathscr{O}_X(p^*D_0)$ for some positive integer m_0 and some divisor D_0 on Z, then $\bar{D}_0$, the closure of D_0 in V, is not ample.*

Proof. Assume the opposite. By replacing m_0 with a large multiple, we may assume that $\bar{D}_0$ is very ample. We may also assume that both $\bar{D}_0$ and p^*D_0 are prime and the pair $(V, \sum_{i=0}^r \frac{m_i-1}{m_i}\bar{D}_i)$ is klt. Let $c_X : X_1 \to X$ be the ramified cyclic cover with respect to $\mathscr{F}$, m_0 and p^*D_0 (see [KM98, Def. 2.52]). Then c_X is ramified over p^*D_0 with degree m and $(c_X^*\mathscr{F})^{**} \cong \mathscr{O}_{X_1}(E)$, where $E = (c_X^*p^*D_0)_{red}$. Moreover, over the smooth locus of X, there is an injection of sheaves from $c_X^*\Omega^1_X$ to $\Omega^1_{X_1}$. Hence by Proposition 2.1, we have an injection $\mathscr{O}_{X_1}(E) \hookrightarrow \Omega^{[1]}_{X_1}$.

Let $X_1 \xrightarrow{p_1} Z_1 \xrightarrow{c_Z} Z$ be the Stein factorisation. Let $c_V : V_1 \to V$ be the normalisation of V in the function field of X_1. Then we obtain an open embedding $j_1 : Z_1 \to V_1$ such that $\mathrm{codim} V_1\backslash Z_1 \geqslant 2$. If F_p is a general fiber of p and if F_{p_1} is a general fiber of p_1 which is mapped to F_p, then $F_{p_1} \to F_p$ is étale. Since p is a Mori fibration, F_p is a smooth rational curve which is simply connected. Hence $F_{p_1} \to F_p$ is an isomorphism and $F_{p_1} \cong \mathbb{P}^1$. Hence c_Z is of degree m_0. Let $H = (c_Z^*D_0)_{red}$. Since $c_Z \circ p_1 : X_1 \to Z$ has connected fibers over D_0, we have $c_Z^*D_0 = m_0H$. Hence $E_1 = p_1^*H$. Note that H is Cartier in codimension 1. By shrinking Z, we may assume that H is Cartier on Z.

Note that $c_X|_{X_1\backslash E}$ is étale in codimension 1 and $c_Z^*D_0 = m_0H$. From Lemma 4.7, we conclude that there is an effective $\mathbb{Q}$-divisor Δ_1' in Z_1 such that $K_{Z_1} + \Delta_1' = p^*(K_Z + \sum_{i=0}^r \frac{m_i-1}{m_i} D_i)$. Hence if Δ_1 is the closure of Δ_1' in V_1, then $K_{V_1} + \Delta_1 = p^*(K_V + \sum_{i=0}^r \frac{m_i-1}{m_i}\bar{D}_i)$. This implies that (V_1, Δ_1) is klt by [KM98, Prop. 5.20 and Cor. 2.35].

Since general fiber of p_1 does not carry any non-zero pluri-forms, by Lemma 2.5, the injection $\mathscr{O}_{X_1}(E_1) \hookrightarrow \Omega^{[1]}_{X_1}$ factorises through $(p_1^*\Omega^1_{Z_1})^{sat}$. Hence by the projection formula, we have an injection from $\mathscr{O}_{Z_1}(H)$ to $p_{1*}((p_1^*\Omega^1_{Z_1})^{sat})$. However, by Lemma 4.6, $p_{1*}((p_1^*\Omega^1_{Z_1})^{sat}) = \Omega^1_{Z_1}$. By Proposition 2.1, we get an injection from $\mathscr{O}_{V_1}(\bar{H})$ to $\Omega^{[1]}_{V_1}$, where $\bar{H}$ is the closure of H in V_1. Hence by Bogomolov-Sommes theorem (see [GKKP11, Thm. 7.2]), the Kodaira dimension of $\mathscr{O}_{V_1}(\bar{H})$ is not larger than 1. However, since $\bar{D}_0$ is ample, $\bar{H}$ is ample and has Kodaira dimension equal to $\dim V_1 = 2$. This leads to a contradiction. □

Lemma 4.9. *Let $p : X \to Z$ be an equidimensional fibration such that general fibers of p do not carry any non-zero pluri-form. Assume that Z is smooth and there is an open embedding $j : Z \to V$ such that* $\mathrm{codim} V\backslash Z \geqslant 2$. *Assume that $H^0(V, \Omega_V^{[r]}) = \{0\}$ for all $r > 0$. Then $H^0(X, \Omega_X^{[r]}) = \{0\}$ for all $r > 0$.*

Proof. Assume the opposite. Let $\mathscr{F} = (p^*\Omega_Z^1)^{sat}$ be the saturation of the image of $p^*\Omega_Z^1$ in $\Omega_X^{[1]}$. Since general fibers of p do not carry any non-zero pluri-form, we have $H^0(X, \Omega_X^{[r]}) \cong H^0(X, \mathscr{F}^{[\wedge r]})$ by Lemma 2.5. Hence there is an injection from $\mathscr{O}_X$ to $\mathscr{F}^{[\wedge r]}$. By taking the direct image, we have an injection from $\mathscr{O}_Z$ to $p_*(\mathscr{F}^{[\wedge r]})$. By Lemma 4.6, $p_*(\mathscr{F}^{[\wedge r]}) \cong \Omega_Z^r$. This implies that $H^0(V, \Omega_V^{[r]}) \neq \{0\}$ by Proposition 2.1, which is a contradiction. □

Now we are ready to prove Theorem 4.2.

Proof of 4.2. We will argue by contradiction. Let $f_Z : Z \to Z'$ be the result of a $(K_Z+\Delta)$-MMP. Assume that Z' has Picard number 1. If $\Delta' = f_{Z*}\Delta$, then $K_{Z'} + \Delta'$ is not pseudo-effective neither. Thus $-(K_{Z'} + \Delta')$ is ample. We know that there is an open subset $Z'_0 \subseteq Z'$ with $\mathrm{codim} Z'\backslash Z'_0 \geqslant 2$ such that f_Z^{-1} is an isomorphism from Z'_0 onto its image. Let Z_0 be $f_Z^{-1}(Z'_0)$, which is an open subset in Z. Since Z' is normal, by shrinking Z'_0 if necessary, we may assume that $Z_0 \cong Z'_0$ are smooth varieties. Let $X_0 = p^{-1}(Z_0)$.

Since $\mathrm{codim} Z'\backslash Z'_0 = 2$, there is a projective curve C'_0 in Z'_0 such that it is an ample divisor in Z'. Let C_0 be the image of C'_0 in Z_0. Let α be a class of movable curves in X whose image in Z is not zero and is proportional to the class of C_0.

Since general fibers of p do not carry any non-zero pluri-forms, by Lemma 2.5, we have $H^0(X, (\Omega_X^1)^{[\otimes m]}) \cong H^0(X, ((p^*\Omega_Z^1)^{sat})^{[\otimes m]})$ for any $m > 0$, where $(p^*\Omega_Z^1)^{sat}$ is the saturation of the image of $(p^*\Omega_Z^1)$ in $\Omega_X^{[1]}$. Hence $\mu_\alpha^{max}((p^*\Omega_Z^1)^{sat}) \geqslant 0$ and there is a coherent sheaf $\mathscr{H}$ saturated in $(p^*\Omega_Z^1)^{sat}$ such that $\mu_\alpha(\mathscr{H}) \geqslant 0$.

However, since $p|_{X_0}$ is a Mori fibration and Z_0 is smooth, the $\mathbb{Q}$-Cartier divisor associates to the determinant of $(p^*\Omega_Z^1)^{sat}|_{X_0}$ is equal to $p^*(K_Z + \Delta)|_{X_0}$ by Remark 4.5. Hence $\det((p^*\Omega_Z^1)^{sat}) \cdot \alpha = p^*(K_Z + \Delta) \cdot \alpha = (p \circ f_Z)^*(K_{Z'} + \Delta') \cdot \alpha < 0$. This implies that $\mathscr{H} \neq (p^*\Omega_Z^1)^{sat}$. If $\mathscr{J}$ is the quotient $(p^*\Omega_Z^1)^{sat}/\mathscr{H}$, then $\mathrm{rank}\,\mathscr{J} = \mathrm{rank}\mathscr{H} = 1$ and $\mathscr{H}[\otimes]\mathscr{J} = \det((p^*\Omega_Z^1)^{sat})$. Thus $\mathscr{J} \cdot \alpha < 0$.

Since $H^0(X, ((p^*\Omega_Z^1)^{sat})^{[\otimes m]}) \neq \{0\}$ for some $m > 0$, we have $H^0(X, \mathscr{H}^{\otimes s}[\otimes]\mathscr{J}^{\otimes t}) \neq \{0\}$ for some $s, t \geqslant 0$ by Lemma 2.3. In this case, we have $s > t$ for $(\mathscr{H}[\otimes]\mathscr{J}) \cdot \alpha < 0$ and $\mathscr{J} \cdot \alpha < 0$.

Let F_p be a general fiber of p. Then the class of F_p is movable and $\mu_{F_p}(\mathscr{H}^{[\otimes s]}[\otimes]\mathscr{J}^{[\otimes t]}) \geqslant 0$. Moreover, we have $(\mathscr{H}[\otimes]\mathscr{J}) \cdot F_p = \det((p^*\Omega_Z^1)^{sat}) \cdot F_p = 0$. This implies that $\mathscr{H} \cdot F_p \geqslant 0$ since $s > t$. However, since the restriction of $(p^*\Omega_Z^1)^{sat}$ on F_p is isomorphic to $\mathscr{O}_{F_p} \oplus \mathscr{O}_{F_p}$, we have $\mathscr{H} \cdot F_p = 0$.

Let k be the smallest positive integer such that $\mathscr{H}^{[\otimes k]}$ is invertible. Then there is a Cartier divisor L in Z such that $\mathscr{H}^{[\otimes k]} \cong \mathscr{O}_X(p^*L)$. Let $L' = f_{Z*}\mathrm{L}$. Then $L' \cdot C'_0 \geqslant 0$ since $\mu_\alpha(\mathscr{H}) \geqslant 0$. Notice that if $L' \cdot C'_0 > 0$, then L' is ample on Z'. Hence by Lemma 4.8, we can only have $L' \cdot C'_0 = 0$ and L' is numerically equal to the zero divisor since Z' has Picard number 1. By [AD12, Lem. 2.6], there is a positive integer k' such that $k'L'$ is linearly equivalent to the zero divisor. Hence $\mathscr{O}_Z(k'L)|_{Z_0} \cong \mathscr{O}_{Z_0}$ and $\mathscr{H}^{[\otimes kk']}|_{X_0} \cong \mathscr{O}_{X_0}$. Let l be the smallest positive integer such that $\mathscr{H}^{[\otimes l]}|_{X_0} \cong \mathscr{O}_{X_0}$. Let $c : W_0 \to X_0$ be the cyclic cover with respect to $\mathscr{H}|_{X_0}$ and l (see [KM98, Def. 2.52]). Then c is étale in codimension 1. We have $\mathscr{O}_{W_0} \cong (c^*\mathscr{H})^{**}$ and there is an injection $(c^*\mathscr{H})^{**} \hookrightarrow (c^*\Omega_{X_0}^1)^{**} \cong \Omega_{W_0}^{[1]}$. Hence $H^0(W_0, \Omega_{W_0}^{[1]}) \neq \{0\}$.

On the other hand, let $W_0 \to V_0 \to Z'_0$ be the Stein factorisation. Let $h : V \to Z'$ be the normalisation of Z' in the function field of W_0. Then there is an open embedding from V_0 to V such that $\mathrm{codim} V\backslash V_0 \geqslant 2$. By Lemma 4.7, we have $K_{V_0} \leqslant (h^*(K_{Z'} + \Delta'))|_{V_0}$. Since $\mathrm{codim} V\backslash V_0 \geqslant 2$, there is an effective $\mathbb{Q}$-divisor Δ_V in V such that $K_V + \Delta_V = h^*(K_{Z'} + \Delta')$. Thus the pair (V, Δ_V) is klt by [KM98, Prop. 5.20]. Moreover, $-(K_V + \Delta_V)$ is ample since $-(K_{Z'} + \Delta')$ is ample, hence V is rationally connected by [HM07, Cor. 1.13]. By Theorem 1.2, we have $H^0(V, \Omega_V^{[1]}) = 0$. However, since general fibers of $W_0 \to V_0$ are isomorphic to $\mathbb{P}^1$ which does not carry any non-zero pluri-form, we conclude that $H^0(W_0, \Omega_{W_0}^{[1]}) = \{0\}$ from Lemma 4.9. This is a contradiction. □

4.2. **The case of threefolds with canonical singularities.** In this subsection, we will study Mori fibrations $X \to Z$ such that X is a projective threefold with $\mathbb{Q}$-factorial canonical singularities and Z is a projective normal surface. Assume that X carries non-zero pluri-forms. Then by Theorem 4.2, either the result Z' of any $(K_Z + \Delta)$-MMP has Picard number 2 or $(K_Z + \Delta)$ is pseudo-effective, where Δ the $\mathbb{Q}$-divisor

defined in Theorem 4.2. We will study the first case in Proposition 4.10 and the second case in Proposition 4.12.

Proposition 4.10. *Let $p : X \to Z$ be a Mori fibration from a projective threefold to a projective surface such that X has at most $\mathbb{Q}$-factorial canonical singularities. Let Δ be the divisor in Z defined in Theorem 4.2. Assume that $K_Z + \Delta$ is not pseudo-effective. Let $f_Z : Z \to Z'$ be the result of a $(K_Z + \Delta)$-MMP and let Δ' be the strict transform of Δ in Z'. Then there is a $(K_{Z'} + \Delta')$-Mori fibration $\pi' : Z' \to \mathbb{P}^1$. Let $\pi = \pi' \circ f_Z$ and $q = \pi \circ p : X \to \mathbb{P}^1$. Then we have $H^0(X, (\Omega^1_X)^{[\otimes m]}) \cong H^0(X, ((q^*\Omega^1_{\mathbb{P}^1})^{sat})^{[\otimes m]})$ for any $m > 0$, where $(q^*\Omega^1_{\mathbb{P}^1})^{sat}$ is the saturation of the image of $q^*\Omega^1_{\mathbb{P}^1}$ in $\Omega^{[1]}_X$.*

Proof. Let $\mathscr{H} = (q^*\Omega^1_{\mathbb{P}^1})^{sat}$ and let $\mathscr{F} = (p^*\Omega^{[1]}_Z)^{sat}$ be the saturation of $p^*\Omega^{[1]}_Z$ in $\Omega^{[1]}_X$. Then we have an exact sequence of coherent sheaves $0 \to \mathscr{H} \to \mathscr{F} \to \mathscr{J} \to 0$, where $\mathscr{J}$ is a torsion free sheaf such that $\det\mathscr{F} = \mathscr{H}[\otimes]\mathscr{J}$. Let α be a class movable curves in X whose image in Z is not zero and is proportional to the class of general fibers of $\pi : Z \to \mathbb{P}^1$. Then $\mathscr{H} \cdot \alpha = 0$. Moreover, we have $(\mathscr{H}[\otimes]\mathscr{J}) \cdot \alpha = \det\mathscr{F} \cdot \alpha = p^*(K_Z + \Delta) \cdot \alpha$ by Remark 4.5. This intersection number is negative since for general fibers F_π of $\pi : \mathrm{Z} \to \mathbb{P}^1$, we have $(K_Z + \Delta) \cdot F_\pi = (K_{Z'} + \Delta') \cdot (f_{Z*}F_\pi) < 0$. Hence $\mathscr{J} \cdot \alpha < 0$.

There is an open subset U of X with $\mathrm{codim}(X \backslash U) \geqslant 2$ such that we have an exact sequence of locally free sheaves over U, $0 \to \mathscr{H}|_U \to \mathscr{F}|_U \to \mathscr{J}|_U \to 0$. Since $\mu_\alpha(\mathscr{H}^{\otimes s} \otimes \mathscr{J}^{\otimes t}) < 0$ if $t > 0$, we have $H^0(U, \mathscr{H}|_U^{\otimes s} \otimes \mathscr{J}|_U^{\otimes t}) = \{0\}$ if $t > 0$. Hence by Lemma 2.5 and Lemma 2.4, we obtain $H^0(X, (\Omega^1_X)^{[\otimes m]}) \cong H^0(X, (\mathscr{F})^{[\otimes m]}) \cong H^0(X, (\mathscr{H})^{[\otimes m]})$ for any $m > 0$. □

Example 4.11. We will give an example of this kind of threefolds. Let $Z = \mathbb{P}^1 \times \mathbb{P}^1$. Denote by pr_1, pr_2 the two natural projections from Z to $\mathbb{P}^1$. Let z_1,...,z_r be $r \geqslant 4$ different points in $\mathbb{P}^1$ and let $C_i = pr_1^* z_i$ for $i = 1, ..., r$. Let $X_0 = \mathbb{P}^1 \times Z$. By the method of Construction 2.13, we can construct a Mori fibration $\pi : X \to Z$ such that $m(\pi, C_i) = 2$ for $i = 1, ..., r$. Notice that $K_Z + \frac{1}{2}(C_1 + \cdots + C_r)$ is not pseudo-effective since it has negative intersection number with general fibers of pr_1. Moreover, we have $H^0(X, (\Omega^1_X)^{[\otimes 2]}) \cong H^0(\mathbb{P}^1, (\Omega^1_{\mathbb{P}^1})^{\otimes 2} \otimes \mathscr{O}_{\mathbb{P}^1}(C_1 + \cdots + C_r)) \cong H^0(\mathbb{P}^1, \mathscr{O}_{\mathbb{P}^1}(-4 + r)) \neq \{0\}$.

Now we will treat the second case. Note that this is the case for Example 2.12 and Example 2.14.

Proposition 4.12. *Let $p : X \to Z$ be a Mori fibration from a projective threefold to a projective surface. Assume that X has at most $\mathbb{Q}$-factorial klt singularities. Assume that $(K_Z + \Delta)$ is pseudo-effective, where Δ is the $\mathbb{Q}$-divisor defined in Theorem 4.2. Then X carries non-zero pluri-forms.*

Proof. By the abundance theorem for log surface (see. [AFKM92]), $(K_Z + \Delta)$ is effective . Hence there is a positive integer l such that $l(K_Z + \Delta)$ is an effective Cartier divisor. This implies that $H^0(X, (\Omega^1_X)^{[\otimes 2l]}) \neq \{0\}$ by the Lemma 4.4. □

5. Proof of Theorem 1.4

In this section, we will complete the proof of Theorem 1.4. First we will show that if X is a rationally connected projective threefold with *terminal* singularities such that $H^0(X, (\Omega^1_X)^{[\otimes m]}) \neq \{0\}$ for some $m > 0$, then there is a dominant rational map from X to $\mathbb{P}^1$ (Lemma 5.2). To this end, we will need the following lemma.

Lemma 5.1. *Let $p : X \to Z$ be a Mori fibration such that X has $\mathbb{Q}$-factorial terminal singularities and $\dim Z = \dim X - 1$. Then there exists an open subset $Z_0 \subseteq Z$ with $\mathrm{codim}(Z \backslash Z_0) \geqslant 2$ such that every scheme-theoretic fiber over Z_0 is reduced.*

Proof. Since X has terminal singularities, it is smooth in codimension 2 (see [KM98, Cor. 5.18]). There is a smooth open subset $Z_0 \subseteq Z$ with $\mathrm{codim}(Z \backslash Z_0) \geqslant 2$ such that $p|_{X_0} : X_0 \to Z_0$ is equidimensional and X_0 is smooth, where $X_0 = q^{-1}(Z_0)$. Since Z_0 is smooth and X is Cohen-Macaulay (see [KM98, Thm. 5.10 and Thm. 5.22]), fibers of q over Z_0 are Cohen-Macaulay. Hence a fiber over Z_0 is reduced if and only if it is generically reduced. If z is a point in Z_0, then there is a smooth curve B in Z_0 passing through z such that $Y = p^{-1}(B)$ is a smooth surface. Moreover, $p|_Y : Y \to B$ is a Fano fibration for p is a Mori fibration. Thus $p|_Y$ has reduced fibers over B by [Ou13, Prop. 3.5]. Hence the scheme-theoretic fiber $p^{-1}(\{z\})$ is reduced. □

Lemma 5.2. *Let X be a rationally connected projective threefold with terminal singularities such that $H^0(X,(\Omega^1_X)^{[\otimes m]}) \neq \{0\}$ for some $m > 0$. Let $f_X : X \dashrightarrow X'$ be the result of a MMP for X. Then there is a fibration $p' : X' \to \mathbb{P}^1$.*

Proof. Notice that X' is a Mori fiber space since K_X is not pseudo-effective by [Kol96, Cor. 1.11]. Then we have a Mori fibration $q' : X' \to Z$, where Z is a normal rationally connected variety. If $\dim Z = 1$, then we are done.

By Lemma 2.2, we know that X' carries non-zero pluri-forms. Hence $\dim Z > 0$ by [Ou13, Thm. 2.1]. Assume that $\dim Z = 2$. Then Z has at most canonical singularities by [MP08, Cor. 1.2.8]. Hence K_Z is not pseudo-effective by [Kol96, Cor. 1.11]. Moreover, by Lemma 5.1, $m(q', D) = 1$ for any effective divisor D on Z. Hence, by Theorem 4.2, if $Z \to Z'$ is the result of a MMP for Z, then Z' has Picard number 2. Hence we have a Mori fibration $Z' \to \mathbb{P}^1$. Let p' be the composition of $X' \to Z \to Z' \to \mathbb{P}^1$. Then p' is a fibration from X' to $\mathbb{P}^1$ □

With the notation above, notice that any general fiber F' of $p' : X' \to \mathbb{P}^1$ is smooth rationally connected surface. Hence F' does not carry any non-zero pluri-form. Let U be the largest open subset in X over which $f_X : X \dashrightarrow X'$ is regular. Then $\mathrm{codim}_X X\backslash U \geqslant 2$, $\mathrm{codim}_{X'} X'\backslash f_X(U) \geqslant 2$ and the rational map $p : X \dashrightarrow \mathbb{P}^1$ is regular over U. If F is a general fiber of $U \to \mathbb{P}^1$, then $f_X(F) \subseteq F'$, where F' is a general fiber of p'. Moreover $\mathrm{codim}_{F'} F'\backslash f_X(F) \geqslant 2$. Hence $f_X(F)$ does not carry any non-zero pluri-form and neither does F by Lemma 2.2. The following lemma shows that the rational map $X \dashrightarrow \mathbb{P}^1$ is regular. Moreover, general fibers of $X \to \mathbb{P}^1$ are birational to the ones of $X' \to p^1$ which are smooth Fano surfaces. Hence general fibers of $X \to \mathbb{P}^1$ are rationally connected.

Lemma 5.3. *Let X be a projective threefold with $\mathbb{Q}$-factorial terminal singularities. Assume that there is a non constant rational map $p : X \dashrightarrow \mathbb{P}^1$ which is regular over U such that $\mathrm{codim} X\backslash U \geqslant 2$. Assume that general fibers of $U \to \mathbb{P}^1$ do not carry any pluri-form. If $H^0(X,(\Omega^1_X)^{[\otimes m]}) \neq \{0\}$ for some $m > 0$, then p is regular.*

Proof. Let Γ be the normalisation of the graph of p. Let $pr_1 : \Gamma \to X$, $pr_2 : \Gamma \to \mathbb{P}^1$ be the natural projections. Then there is a natural injection from $H^0(\Gamma,(\Omega^1_\Gamma)^{[\otimes m]})$ to $H^0(X,(\Omega^1_X)^{[\otimes m]})$ by Lemma 2.2. Let $\sigma \in H^0(X,(\Omega^1_X)^{[\otimes m]})$ be a non-zero element. Then σ induces a rational section σ_Γ of $(\Omega^1_\Gamma)^{[\otimes m]}$ on Γ. Let E be a pr_1-exceptional divisor. Then there is a curve in E which is contracted by pr_2 since $\dim E = 2 > 1$. Hence this curve is not contracted by pr_1 since the graph of p is included in $X \times \mathbb{P}^1$ and the normalisation map is finite. Thus $pr_1(E)$ is a curve in X and X is smooth around the generic point of $pr_1(E)$ since X is smooth in codimension 2 (see [KM98, Cor. 5.18]). Hence σ_Γ does not have pole along E. This implies that we have an isomorphism from $H^0(\Gamma,(\Omega^1_\Gamma)^{[\otimes m]})$ to $H^0(X,(\Omega^1_X)^{[\otimes m]})$ induced by pr_1.

Note that $pr_1^{-1}|_U$ induces an isomorphism from U onto its image. If F_U is a general fiber of $p|_U$, then $pr_1^{-1}(F_U)$ is an open subset of F_Γ, where F_Γ is a general fiber of $pr_2 : \Gamma \to \mathbb{P}^1$. Since F_U does not carry any non-zero pluri-form, neither does F_Γ. Hence by Lemma 2.5, we have $H^0(\Gamma,(\Omega^1_\Gamma)^{[\otimes m]}) \cong H^0(\Gamma,((pr_2^*\Omega^1_{\mathbb{P}^1})^{sat})^{[\otimes m]})$.

We will first prove that p is regular in codimension 2. Assume the opposite. Then there is a divisor D in Γ which is exceptional for $pr_1 : \Gamma \to X$ and $\mathrm{codim}_X pr_1(D) = 2$. Since X is smooth in codimension 2 (see [KM98, Cor. 5.18]), X is smooth around the generic point of $pr_1(D)$. Thus there is a smooth quasi-projective curve C in D such that Γ is smooth along C and $pr_1(C)$ is a smooth point in X. Notice that C is horizontal over $\mathbb{P}^1$ under the projection $pr_2 : \Gamma \to \mathbb{P}^1$ for the same reason as above. Let σ be a non-zero element in $H^0(\Gamma,(\Omega^1_\Gamma)^{[\otimes m]})$. By the exact sequence of locally free sheaves $\Omega^1_\Gamma|_C \to \Omega^1_C \to 0$, σ induce an element σ_C in $H^0(C,(\Omega^1_C)^{\otimes m})$. On one hand, C is horizontal over $\mathbb{P}^1$ and σ is non-zero in $H^0(\Gamma,((pr_2^*\Omega^1_{\mathbb{P}^1})^{sat})^{[\otimes m]})$, we have $\sigma_C \neq 0$. On the other hand, $pr_1(C)$ is a smooth point in X. Hence we obtain $\sigma_C = 0$ since σ is the pullback of certain element in $H^0(X,(\Omega^1_X)^{[\otimes m]})$. This is a contradiction.

Now we will prove that p is regular. Let F_1 and F_2 be two different fibers of $U \to \mathbb{P}^1$. Then their closures in X are two Weil divisors and their intersection is included in a closed subset of codimension at most 3. Hence their intersection is empty since X is $\mathbb{Q}$-factorial. This implies that p is regular. □

Together with Lemma 2.5 and Lemma 3.1, we can conclude Theorem 1.4.

References

[AD12] Carolina Araujo and Stéphane Druel, *On codimension 1 del pezzo foliations on varieties with mild singularities*, Preprint `arXi:1210.4013`, 2012.

[AFKM92] Dan Abramovich, L.-Y. Fong, János Kollár, and James Mckernan, *Semi log canonical surface*, Astérisque **211** (1992), 139–154.

[Cam92] Frédéric Campana, *Connexité rationnelle des variétés de fano*, Annales scientifiques de l'Ecole normale supérieure, vol. 25, Société mathématique de France, 1992, pp. 539–545.

[Fuj99] Osamu Fujino, *Applications of Kawamata's positivity theorem*, Proc. Japan Acad. Ser. A Math. Sci. **75** (1999), no. 6, 75–79. MR 1712648 (2000f:14089)

[GHS03] Tom Graber, Joe Harris, and Jason Starr, *Families of rationally connected varieties*, Journal of the American Mathematical Society **16** (2003), no. 1, 57–67.

[GKKP11] Daniel Greb, Stefan Kebekus, Sándor J. Kovács, and Thomas Peternell, *Differential forms on log canonical spaces*, Publ. Math. Inst. Hautes Études Sci. (2011), no. 114, 87–169. MR 2854859

[GKP12] Daniel Greb, Stefan Kebekus, and Thomas Peternell, *Reflexive differential forms on singular spaces – geometry and cohomology*, Preprint `arXiv: arXiv:1202.3243v2`, 2012.

[Har77] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977, Graduate Texts in Mathematics, No. 52.

[Har80] ———, *Stable reflexive sheaves*, Math. Ann. **254** (1980), no. 2, 121–176. MR 597077 (82b:14011)

[HM07] Christopher D. Hacon and James Mckernan, *On Shokurov's rational connectedness conjecture*, Duke Math. J. **138** (2007), no. 1, 119–136. MR 2309156 (2008f:14030)

[Kaw92] Yujiro Kawamata, *Termination of log flips for algebraic 3-folds*, Internat. J. Math **3** (1992), no. 5, 653–659.

[KM98] János Kollár and Shigefumi Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998, With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original. MR 1658959 (2000b:14018)

[KMM87] Yujiro Kawamata, Katsumi Matsuda, and Kenji Matsuki, *Introduction to the minimal model problem*, Algebraic geometry, Sendai, 1985, Adv. Stud. Pure Math., vol. 10, North-Holland, Amsterdam, 1987, pp. 283–360. MR 946243 (89e:14015)

[KO82] Shoshichi Kobayashi and Takushiro Ochiai, *Holomorphic structures modeled after hyperquadrics*, Tohoku Mathematical Journal **34** (1982), no. 4, 587–629.

[Kol96] János Kollár, *Rational curves on algebraic varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 32, Springer-Verlag, Berlin, 1996. MR 1440180 (98c:14001)

[Kol08] János Kollár, *Is there a topological Bogomolov-Miyaoka-Yau inequality?*, Pure Appl. Math. Q. **4** (2008), no. 2, 203–236.

[Laz04] Robert Lazarsfeld, *Positivity in algebraic geometry. I*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 48, Springer-Verlag, Berlin, 2004, Classical setting: line bundles and linear series. MR 2095471 (2005k:14001a)

[Mor88] Shigefumi Mori, *Flip theorem and the existence of minimal models for 3-folds*, Journal of the American Mathematical Society **1** (1988), no. 1, 117–253.

[MP97] Yoichi Miyaoka and Thomas Peternell, *Geometry of higher-dimensional algebraic varieties*, DMV Seminar, vol. 26, Birkhäuser Verlag, Basel, 1997. MR 1468476 (98g:14001)

[MP08] Shigefumi Mori and Yuri Prokhorov, *On $\mathbb{Q}$-conic bundles*, Publ. Res. Inst. Math. Sci. **44** (2008), no. 2, 315–369. MR 2426350 (2009e:14062)

[Ou13] Wenhao Ou, *Singular rationally connected surfaces with non-zero pluri-forms*, Preprint `arXiv:1301.0915`, 2013.

[Rei79] Miles Reid, *Canonical 3-folds*, Journées de géométrie algébrique dAngers (1979), 273–310.

[Rei87] ———, *Young persons guide to canonical singularities*, Proc. Symp. Pure Math, vol. 46, 1987, pp. 345–414.

[Tot10] Burt Totaro, *Algebraic surfaces and hyperbolic geometry*, Preprint `arXiv:1008.3825`, 2010.

Wenhao OU: Institut Fourier, UMR 5582 du CNRS, Université Grenoble 1, BP 74, 38402 Saint Martin d'Hères, France